


【Claims】

【claim 1】

5 A compound of the general formula(I)



wherein Y is zero or



wherein X is oxygen or sulfur, R₁, R₂, R₃, R₄ and R₅ are independently hydrogen, halogen, nitro, amino, hydroxy, C₁-C₄ lower alkylhydroxy, C₁-C₄ lower alkylamino, C₁-C₈ alkyl or C₁-C₄ lower alkoxy, R' and R'' are independently C₁-C₈ alkyl or C₁-C₄ lower alkoxy, and Z is C₁-C₄ lower alkyl, C₁-C₄ lower alkoxy or C₁-C₄ lower alkylamino or pharmaceutically acceptable salt thereof.



25

30


- 34 -

[claim 2]

A process for the preparation of a compound of the following general formula (I) or pharmaceutically acceptable salt thereof, comprising reacting a compound of the following general formula(a) with a compound of the following general formula(b) to give a compound of the following general formula (I) and if necessary converting the compound of the general formula (I) into pharmaceutically acceptable salt thereof.




wherein R_1 , R_2 , R_3 , R_4 , R_5 , R' , R'' , X , Y and Z are as defined above and




- 35 -

[claim 3]


A process for the preparation of a compound of the following general formula (I) or pharmaceutically acceptable salt thereof, comprising reacting a compound of the following general formula(c) with a compound of the following general formula(d) to give a compound of the following general formula (I) and if necessary converting the compound of the general formula (I) into pharmaceutically acceptable salt thereof.



(I)



(a)



(b)

wherein R_1 , R_2 , R_3 , R_4 , R_5 , R' , R'' , X , Y and Z are as defined above and

